

Meta-Analysis in R with the *metafor* Package

Evidence Synthesis Hackathon
Stockholm Environment Institute

Wolfgang Viechtbauer

Maastricht University

2018-04-23

A Bit of History

- research focused on the statistical methods for meta-analysis
- there were several R packages for meta-analysis in 2000, but all lacked 'meta-regression' capabilities
- wrote a function for this and put it on my website
- turned this into a full R package (**metafor**) in 2010
- metafor: "meta-analysis package for R" (easier than: metapforR)

1

2

Example: BCG Vaccine

- BCG: Bacillus Calmette-Guérin
- BCG is a vaccine against tuberculosis
- effectiveness studies: compare proportion of TB positive cases in vaccinated and non-vaccinated group

Results from One Trial

	Positive	Negative	Total
Vaccinated	4	119	123
Not Vaccinated	11	128	139

$$RR = \frac{4/123}{11/139} = .41$$
$$p_T = 4/123 = .0325$$
$$p_C = 11/139 = .0791$$
$$y = \ln[RR] = \ln \left[\frac{4/123}{11/139} \right] = -.89$$
$$v = \frac{1}{4} - \frac{1}{123} + \frac{1}{11} - \frac{1}{139} = .326$$

3

4

Results from 13 Trials

```
> library(metafor)
> dat.bcg

##   trial      author year tpos  tneg cpos  cneg ablat    alloc
## 1     1       Aronson 1948   4   119   11  128   44  random
## 2     2 Ferguson & Simes 1949   6   300   29  274   55  random
## 3     3 Rosenthal et al 1960   3   228   11  209   42  random
## 4     4 Hart & Sutherland 1977  62 13536  248 12619  52  random
## 5     5 Frimodt-Møller et al 1973  33  5036   47  5761  13 alternate
## 6     6 Stein & Aronson 1953  180 1361   372 1079   44 alternate
## 7     7 Vandiviere et al 1973   8  2537   10  619   19  random
## 8     8 TPT Madras 1980  505 87886  499 87892  13  random
## 9     9 Coetze & Berjak 1968   29  7470   45  7232  27  random
## 10   10 Rosenthal et al 1961  17  1699   65  1600  42 systematic
## 11   11 Comstock et al 1974  186 50448  141 27197  18 systematic
## 12   12 Comstock & Webster 1969   5  2493   3  2338  33 systematic
## 13   13 Comstock et al 1976  27 16886  29 17825  33 systematic
```

Calculate Outcome Measure

```
> dat <- escalc(measure="RR", ai=tpos, bi=tneg, ci=cpos, di=cneg, data=dat.bcg)
> dat

##   trial      author year ... yi    vi
## 1     1       Aronson 1948 ... -0.8893 0.3256
## 2     2 Ferguson & Simes 1949 ... -1.5854 0.1946
## 3     3 Rosenthal et al 1960 ... -1.3481 0.4154
## 4     4 Hart & Sutherland 1977 ... -1.4416 0.0200
## 5     5 Frimodt-Møller et al 1973 ... -0.2175 0.0512
## 6     6 Stein & Aronson 1953 ... -0.7861 0.0069
## 7     7 Vandiviere et al 1973 ... -1.6209 0.2230
## 8     8 TPT Madras 1980 ... 0.0120 0.0040
## 9     9 Coetze & Berjak 1968 ... -0.4694 0.0564
## 10   10 Rosenthal et al 1961 ... -1.3713 0.0730
## 11   11 Comstock et al 1974 ... -0.3394 0.0124
## 12   12 Comstock & Webster 1969 ... 0.4459 0.5325
## 13   13 Comstock et al 1976 ... -0.0173 0.0714
```

5


6

Meta-Analysis

```
> rma(yi, vi, data=dat)


##
## Random-Effects Model (k = 13; tau^2 estimator: REML)
##
## tau^2 (estimated amount of total heterogeneity): 0.3132 (SE = 0.1664)
## tau (square root of estimated tau^2 value):      0.5597
## I^2 (total heterogeneity / total variability):   92.22%
## H^2 (total variability / sampling variability): 12.86
##
## Test for Heterogeneity:
## Q(df = 12) = 152.2330, p-val < .0001
##
## Model Results:
##
## estimate    se    zval   pval ci.lb ci.ub
## -0.7145  0.1798 -3.9744 <.0001 -1.0669 -0.3622 ***
## ...
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Forest Plot



8

Potential Moderators?




Moderator Analysis

```
> rma(yi, vi, mods = ~ ablat + year, data=dat)


##
## Mixed-Effects Model (k = 13; tau^2 estimator: REML)
##
## tau^2 (estimated amount of residual heterogeneity):  0.1108 (SE = 0.0845)
## tau (square root of estimated tau^2 value):        0.3328
## I^2 (residual heterogeneity / unaccounted variability): 71.98%
## H^2 (unaccounted variability / sampling variability): 3.57
## R^2 (amount of heterogeneity accounted for):       64.63%
##
## Test for Residual Heterogeneity:
## QE(df = 10) = 28.3251, p-val = 0.0016
##
## Test of Moderators (coefficients 2-3):
## QM(df = 2) = 12.2043, p-val = 0.0022
##
## Model Results:
##
## estimate    se    zval   pval ci.lb ci.ub
## intcpt  29.0959  0.9838 -0.1219  0.9883 -60.5724 53.4814
## ablat   -0.8280  0.0102 -2.7371  0.0062 -0.0481 -0.0888 **
## year     0.0019  0.0147  0.1299  0.8966 -0.0269  0.0307
## ...
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10

Moderator Analysis




Forest Plot with Subgroups




11

12

Funnel Plots



Cumulative Meta-Analysis



Models and Analysis Approaches

- fixed-, random-, and mixed-effects models (`rma()`)
- Mantel-Haenszel and Peto's method for 2x2 table and two-group person-time data (`rma.mh()`, `rma.peto()`)
- generalized linear models (i.e., mixed-effects logistic and Poisson regression models) for 2x2 table data, two-group person-time data, proportions, and incidence rates (`rma.glmm()`)
- multilevel and multivariate meta-analysis (`rma.mv()`)
- network meta-analysis (`rma.mv()`)
- phylogenetic meta-analysis (`rma.mv()`)
- spatio-temporal models (`rma.mv()`)

Plots

- forest plots (`forest()`)
- funnel plots (`funnel()`)
- radial (Galbraith) plots (`radial()`)
- Baujat plots (`baujat()`)
- GOSH plots (`gosh()`)
- L'Abbe plots (`labbe()`)

Publication Bias

- rank correlation test (`ranktest()`)
- Egger's regression test (`regtest()`)
- trim and fill method (`trimfill()`)
- Henmi and Copas method (`hc()`)
- file drawer analysis (`fsn()`)

Inference Methods

- fitted / predicted values (`fitted()`, `predict()`)
- best linear unbiased predictions (`ranef()`, `blup()`)
- tests / confidence intervals using the Knapp & Hartung method
- permutation tests/confidence intervals (`permuteest()`)
- likelihood ratio tests (`anova()`)
- profile likelihood confidence intervals (`confint()`)
- (cluster) robust tests / confidence intervals (`robust()`)
- model fit criteria (`logLik()`, `deviance()`)
- information criteria (`AIC()`, `BIC()`, `fitstats()`)
- cumulative meta-analysis (`cumul()`)

Outlier and Influence Diagnostics

- raw, standardized, and studentized residuals (`residuals()`,
`rstandard()`, `rstudent()`)
- DFFITS values, Cook's distances, covariance ratios, and DFBETAS
values (`influence()`)
- leave-one-out analyses (`leave1out()`)
- model weights / hat values (`weights()`, `hatvalues()`)

References / Links

- Viechtbauer, W. (2010). Conducting meta-analyses in R with the
metafor package. *Journal of Statistical Software*, 36(3), 1–48.
- <http://www.metafor-project.org/>
- <https://github.com/wviechtb/metafor>